

Vacunas con marca propia

BQ-BOT is a vaccine for protection of cattle, sheep and goats against Botulism types C and D and Blackquarter. Contains inactivated alum precipitated toxoids obtained from cultures in liquid media of highly toxigenic strains of *Clostridium botulinum* types C and D, and anaculture of *Clostridium chauvoei*.

Dosage: cattle, sheep and goats: 2 ml

Administration: Inject by subcutaneous route. Cattle should first be vaccinated at 6 months and sheep and goats at 4 months of age and annually thereafter. For animals that have never been immunized before, two doses should be given separated by an interval of 4 - 6 weeks.

For veterinary use

Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



bq-bot

Botulism and Blackquarter
Vaccine

100ml

Precautions: Shake well before use. If hypersensitivity reaction develops, adrenaline and / or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 7 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable.

Registration No. G 2549 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01082SA - Expiry date:

PRONDISTAR is a vaccine for protection of cattle, sheep and goats against Anthrax, Botulism types C and D and Blackquarter. Contains inactivated alum precipitated toxoids obtained from cultures in liquid media of highly toxigenic strains of *Clostridium botulinum* types C and D, anaculture of *Clostridium chauvoei*, and not less than 10 million viable spores per dose of an uncapsulated avirulent strain (Sterne 34F2) of *Bacillus anthracis*.

Dosage: cattle, sheep and goats: 2 ml

Administration: Inject by subcutaneous route. Cattle should first be vaccinated at 6 months and sheep and goats at 4 months of age and annually thereafter. For animals that have never been immunized before, two doses should be given separated by an interval of 4 - 6 weeks.

For veterinary use

Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



prondistar

Anthrax, Blackquarter and
Botulism Vaccine

100ml

Precautions: Shake well before use. Pregnant animals can be vaccinated but precautions have to be observed specially in late pregnancy. Seek for professional advice. Antibiotics should not be given for several days before and at least 15 days after vaccination. If hypersensitivity reaction develops, adrenaline and / or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 14 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable. Leftover vaccine, empty vials, and equipment used for vaccinating are contaminated with the live spores and should be autoclaved, disinfected, or incinerated.

Registration No. G 2640 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01092SA - Expiry date:

PRONDITHRAX is a spore vaccine for protection of domestic animals against Anthrax. The vaccine consists of a suspension of not less than 10 million viable spores per ml of an uncapsulated avirulent strain (Sterne 34F2) of *Bacillus anthracis*. Contains saponin as adjuvant.

Dosage: cattle, camels and horses: 1ml
sheep, goats and pigs: 0.5ml

Administration: Inject by subcutaneous route. Animals can be vaccinated from 3 months of age in single doses and annually thereafter. Horses may be slow to develop immunity and consequently a two-dose initial vaccination, administered 1 month apart, followed by a single annual booster is advisable. In very high risk areas, vaccination each six months is recommended.

For veterinary use

Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



prondithrax

Anthrax Spore
Vaccine

100ml

Precautions: Shake well before use. Pregnant animals can be vaccinated but precautions have to be observed specially in late pregnancy. Seek for professional advice. Antibiotics should not be given for several days before and at least 15 days after vaccination. If hypersensitivity reaction develops, adrenaline and/or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 14 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable. Leftover vaccine, empty vials, and equipment used for vaccinating are contaminated with the live spores and should be autoclaved, disinfected, or incinerated.

Registration No. G 2545 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01062SA - Expiry date:

Contains: Toxoids of *Cl.septicum*, *Cl.novyi* type B, *Cl.perfringens* type C, *Cl.perfringens* type D, *Cl.tetani*, and anaculture of *Cl.chauvoei*.

Adjuvant: Alum

Dosage: Cattle: 4 ml
Sheep: 2 ml

Inject by subcutaneous route
Withdrawal period: Nil

Read enclosed package insert for full directions

Batch:
Expiry date:



prondiclos
Polyvalent Clostridial
Vaccine

Cattle, Sheep 100ml

An inactivated vaccine for protection of cattle and sheep against blackleg, enterotoxaemias, gas gangrenes and tetanus.

For veterinary use

Shake well
Store between +2°C and +8°C
Do not freeze
Protect from light

Prondil S.A.
Phone: (598-2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402
Montevideo, URUGUAY

Lot: 01052SC

BQ-VAX is a vaccine for protection of cattle, sheep and goats against Blackquarter. Contains inactivated alum precipitated anaculture of *Clostridium chauvoei*.

Dosage: cattle: 2 ml
sheep and goats: 1 ml

Administration: Inject by subcutaneous route. Cattle should first be vaccinated at 6 months and sheep and goats at 4 months of age and annually thereafter. For animals that have never been immunized before, two doses should be given separated by an interval of 4 - 6 weeks. It is a sound practice to reinoculate vaccinated sheep shortly before the period of greatest risk, specially before any operation likely to produce wounds (lambling, docking, castration, shearing).

For veterinary use
Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



bq-vax

Blackquarter Vaccine

100ml

Precautions: Shake well before use. If hypersensitivity reaction develops, adrenaline and / or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 7 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable.

Registration No. G 2550 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01112SA - Expiry date:

PRONDIBOT is a vaccine for protection of cattle, horses, sheep and goats against Botulism types C and D. Contains inactivated alum precipitated toxoids obtained from cultures in liquid media of highly toxigenic strains of *Clostridium botulinum* types C and D.

Dosage: cattle and horses: 2 ml
sheep and goats: 1 ml

Administration: Inject by subcutaneous route. Cattle and horses should first be vaccinated at 6 months and sheep and goats at 4 months of age and annually thereafter. For animals that have never been immunized before, two doses should be given separated by an interval of 4 - 6 weeks.

For veterinary use
Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



prondibot

Botulism Vaccine

100ml

Precautions: Shake well before use. Pregnant animals can be vaccinated but precautions have to be observed specially in late pregnancy. Seek for professional advice. If hypersensitivity reaction develops, adrenaline and / or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 7 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable.

Registration No. G 2546 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01072SA - Expiry date:

BQ-THRAX is a vaccine for protection of cattle, sheep and goats against Anthrax and Blackquarter. Contains inactivated alum precipitated anaculture of *Clostridium chauvoei*, and not less than 10 million viable spores per dose of an uncapsulated avirulent strain (Sterne 34F2) of *Bacillus anthracis*.

Dosage: cattle, sheep and goats: 2 ml

Administration: Inject by subcutaneous route. Cattle should first be vaccinated at 6 months and sheep and goats at 4 months of age and annually thereafter. For animals that have never been immunized before, two doses should be given separated by an interval of 4 - 6 weeks.

For veterinary use
Storage: Store between +2°C and +8°C. Do not freeze. Protect from light.



bq-thrax

Anthrax and Blackquarter
Vaccine

100ml

Precautions: Shake well before use. Pregnant animals can be vaccinated but precautions have to be observed specially in late pregnancy. Seek for professional advice. Antibiotics should not be given for several days before and at least 15 days after vaccination. If hypersensitivity reaction develops, adrenaline and / or antihistamines should be administered without delay. Animals should not be slaughtered for human consumption within 14 days of vaccination. Always consult a veterinarian prior to vaccinating goats. Development of immunity in goats is variable. Leftover vaccine, empty vials, and equipment used for vaccinating are contaminated with the live spores and should be autoclaved, disinfected, or incinerated.

Registration No. G 2547 (South Africa)

Prondil S.A.
Phone: (598 2) 513 32 54
E-mail: pron@netgate.com.uy
Barros Arana 5402, Montevideo - URUGUAY

Lot: 01102SA - Expiry date: